

Liver Cells Proliferation and Apoptosis in Patients with Alcoholic Liver Disease After Autologous Hematopoietic Stem Cell Transplantation

Burganova G., Abdulkhakov S., Gazizov I., Gumerova A., Titova M., Odintcova A., Kiassov A.
Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2016, Springer Science+Business Media New York. Alcoholic liver disease is a huge medical and social problem that leads to liver fibrosis, cirrhosis, and hepatocellular carcinoma. Unfortunately, the last stages of the disease do not have efficient treatment, except liver transplantation, and require development of new therapeutical approaches. Transplantation of stem cells might be the most promising approach. In our research, we studied transplantation of autologous hematopoietic stem cells (HSC) into the celiac trunk of patients with alcoholic liver cirrhosis. In this article, we pay particular attention to proliferation and apoptosis—two fundamental processes, which determine the fate of regeneration. Liver biopsy specimens before treatment, 3 and 12 months after transplantation of HSC, were stained immunohistochemically with antibodies against PCNA and Bcl-2. The results showed that treatment was safe and effective, hepatocytes increased proliferation, and inflammatory cells decreased antiapoptotic activity, signifying improvement in liver regeneration. However, effect of treatment after 12 months decreases and requires repeated HSC transplantation.

<http://dx.doi.org/10.1007/s12668-016-0354-2>

Keywords

Alcoholic liver disease, Hematopoietic stem cells, Liver cirrhosis, Liver fibrosis, Regeneration, Transplantation

References

- [1] Fujii, H., Hirose, T., Oe, S., et al. (2002). Contribution of bone marrow cells to liver regeneration after partial hepatectomy in mice. *J Hepatol*, 36(5), 653–659.
- [2] Masson, S., Harrison, D. J., Plevvris, J. N., Newsome, P. H. (2004). Potential of hematopoietic stem cell therapy in hepatology: a critical review. *Stem cells*, 22(6), 897–907.
- [3] Kakinuma, S., Tanaka, Y., Chinzei, R., et al. (2003). Human umbilical cord blood as a source of transplantable hepatic progenitor cells. *Stem Cells*, 21(2), 217–227.
- [4] Tang, X. P., Yang, X., Tan, H., Ding, Y. L., Zhang, M., Wang, W. L. (2003). Clinical and experimental study on therapeutic effect of umbilical cord blood transplantation on severe viral hepatitis. *World J. Gastroenterol*, 9(9), 1999–2003.
- [5] Am Esch, J. S., Il, Knoefel, W. T., Klein, M., et al. (2005). Portal application of autologous CD133+ bone marrow cells to the liver: a novel concept to support hepatic regeneration. *Stem Cells*, 23(4), 463–470.

- [6] Am Esch, J. S., II, Schmelzle, M., Fürst, G., et al. (2012). Infusion of CD133+ bone marrow-derived stem cells after selective portal vein embolization enhances functional hepatic reserves after extended right hepatectomy: a retrospective single-center study. *Ann. Surg*, 255(1), 79-85.
- [7] Saadeh, S., Cammel, G., Carey, W. D., Younossi, Z., Barnes, D., Easley, K. (2001). The role of liver biopsy in chronic hepatitis C. *Hepatology*, 33(1), 196-200.
- [8] Leonardi, E., Girlando, S., Serio, G., Mauri, F. A., Perrone, G., Scampini, S., et al. (1992). PCNA and Ki67 expression in breast carcinoma: correlations with clinical and biological variables. *J Clin. Pathol*, 45(5), 416-419.
- [9] Vaux, D. L., Cory, S., Adams, J. M. (1988). Bcl-2 gene promotes haemopoietic cell survival and cooperates with c-myc to immortalize pre-B cells. *Nature*, 335, 440-442.
- [10] Kim, J. K., Park, Y. N., Kim, J. S., et al. (2010). Autologous bone marrow infusion activates the progenitor cell compartment in patients with advanced liver cirrhosis. *Cell Transplant*, 19(10), 1237-1246.
- [11] Terai, S., Sakaida, I., Yamamoto, N., et al. (2003). An in vivo model for monitoring transdifferentiation of bone marrow cells into functional hepatocytes. *J. Biochem*, 134(4), 551-558.